

AD-A161 435

ESTIMATION OF STEADY-STATE CENTRAL MOMENTS BY THE
REGENERATIVE METHOD OF SIMULATION(U) STANFORD UNIV CA
SYSTEMS OPTIMIZATION LAB P W GLYNN ET AL AUG 85

1/1


UNCLASSIFIED

ARO-20927 11-MA DAAG29-84-K-0030

F/G 12/1

NL

							END						
							FINED						
							DATA						



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

②

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD-A161 435

DTIC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ARO 20927.11-MA	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ESTIMATION OF STEADY-STATE CENTRAL MOMENTS BY THE REGENERATIVE METHOD OF SIMULATION		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Peter W. Glynn and Donald L. Iglehart		8. CONTRACT OR GRANT NUMBER(s) DAAG29-84-K-0030
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Operations Research - SOL Stanford University Stanford, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE August 1985
		13. NUMBER OF PAGES p. 16
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) #767 statistical analysis of simulation output #561 regenerative processes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let X be a positive recurrent regenerative process on state space S with steady-state distribution π . Given a function $f : S \rightarrow R$, we consider the problem of estimating the steady-state central moments $\mu_k(f) = \int_S (f(x)-r)^k \pi(dx)$ where r is the steady-state mean of $f(X(\cdot))$. We obtain strong laws, central limit theorems, and confidence intervals for our estimators, and present numerical results.		

DTIC
ELECTE
NOV 22 1985
S
E

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

85 11 18 253

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ESTIMATION OF STEADY-STATE CENTRAL MOMENTS BY THE
REGENERATIVE METHOD OF SIMULATION

by

Peter W. Glynn

and

Donald L. Iglehart

TECHNICAL REPORT NO. 11

August 1985

Prepared under the Auspices
of

U.S. Army Research Contract*
DAAG29-84-K-0030

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Approved for public release: distribution unlimited.

Reproduction in Whole or in Part is Permitted for any
Purpose of the United States Government

DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

* This research was also partially supported under
National Science Foundation Grant MCS-8203483.

ABSTRACT

Let X be a positive recurrent regenerative process on state space S with steady-state distribution π . Given a function $f : S \rightarrow \mathbb{R}$, we consider the problem of estimating the steady-state central moments $\mu_k(f) = \int_S (f(x) - r)^k \pi(dx)$ where r is the steady-state mean of $f(X(\cdot))$. We obtain strong laws, central limit theorems, and confidence intervals for our estimators, and present numerical results.

KEYWORDS:

#767 statistical analysis of simulation output

#561 regenerative processes

1. INTRODUCTION

Let $X = \{X(t) : t \leq 0\}$ be a (possibly) delayed regenerative process taking values in state space S , with regeneration times $T(-1) = 0 \leq T(0) < T(1) < \dots$ (to incorporate regenerative sequences $\{X_n : n \geq 0\}$, we pass to the continuous time process $X(\cdot)$ defined by $X(t) = X_{[t]}$, where $[t]$ is the greatest integer less than or equal to t). Under quite general conditions (see, for example, p. 185 of HEYMAN and SOBEL (1982)), there exists a probability distribution π on S such that

$$(1.1) \quad r_t(f) = \frac{1}{t} \int_0^t f(X(s)) ds \rightarrow \int_S f(y) \pi(dy) \equiv r(f)$$

a.s. as $t \rightarrow \infty$, for a broad class of functions $f: S \rightarrow \mathbb{R}$.

An important problem that has been extensively studied in the simulation literature concerns the estimation of the parameter $r(f)$; $r(f)$ has the interpretation, as is clear from (1.1), as the steady-state mean of $f(X(\cdot))$. In certain applications, however, it may also be of interest to estimate the fluctuations of $f(X(\cdot))$ around its steady-state limit. To be precise, set $f_c(\cdot) = f(\cdot) - r(f)$ and put $v(f) = r(f_c^2)$ (for $g: S \rightarrow \mathbb{R}$, define $g^m: S \rightarrow \mathbb{R}$ via $g^m(x) = g(x) \cdot g(x) \cdots g(x)$ (m times)). Letting f_c^2 play the role of f in (1.1), we observe that $v(f)$ may be regarded as the steady-state variance of $f(X(\cdot))$.

More generally, let $\mu_m(f) = r(f_c^m)$; then $\mu_m(f)$ is the m 'th steady-state central moment of $f(\cdot)$. Clearly, $\mu_1(f) = 0$ and $\mu_2(f) = v(f)$. Note that estimation of $\mu_m(f) = r(f_c^m)$ is not a special case of the standard regenerative method (see, for example, IGLEHART (1978)), since f_c depends on the unknown parameter $r(f)$, which itself must be

estimated. Our goal, in this paper, is to develop an estimation methodology for the central moments, $\mu_m(f)$. In Section 2, we prove the required limit theorems upon which our methods are based. Section 3 develops confidence intervals for central moments, and numerical results are presented in Section 4.

2. ESTIMATORS AND LIMIT THEOREMS FOR CENTRAL MOMENTS

We assume throughout the remainder of this paper that:

$$(2.1) \quad E(Y_1(f)^{2m} + \tau_1^{2m}) < \infty$$

where

$$\tau_n = T(n) - T(n-1) \quad \text{and} \quad Y_n(g) = \int_{T(n-1)}^{T(n)} g(X(s)) ds$$

for functions $g : S \rightarrow R$. Our goal is to estimate the central moments $\mu_k(f)$, $1 \leq k \leq m$.

The following binomial representation of the central moments is crucial to our development:

$$(2.2) \quad \begin{aligned} \mu_k(f) &= r((f - r(f)))^k \\ &= \sum_{j=0}^k \binom{k}{j} r(f^j) (-1)^{k-j} r(f)^{k-j} . \end{aligned}$$

For $g : S \rightarrow R$, set $r(n, g) = \bar{Y}_n(g) / \bar{\tau}_n$, where

$$\bar{Y}_n(g) = \frac{1}{n} \sum_{k=1}^n Y_k(g)$$

and

$$\bar{\tau}_n = \frac{1}{n} \sum_{k=1}^n \tau_k.$$

Relation (2.2) suggests that

$$(2.3) \quad u(n,k) = \sum_{j=0}^k \binom{k}{j} r(n, f^j) (-1)^{k-j} r(n, f)^{k-j}$$

should be a reasonable estimator for $\mu_k(f)$.

(2.4) PROPOSITION. $u(n,k) \rightarrow \mu_k(f)$ a.s. as $n \rightarrow \infty$, for $1 \leq k \leq 2m$.

PROOF. From (2.2) and (2.3), it is clear that we need only show that $r(n, f^k) \rightarrow r(f^k)$ a.s. as $n \rightarrow \infty$, for $k \leq 2m$. The strong law of large numbers applies to both the numerator $\bar{Y}_n(f^k)$ and denominator $\bar{\tau}_n$, yielding the required convergence, provided that $E|Y_1(f^k)| < \infty$ and $E\tau_1 < \infty$. Clearly, (2.1) implies that $E\tau_1 < \infty$, whereas the inequality

$$\begin{aligned} E|Y_1(f^k)| &\leq E\left(\int_{T(0)}^{T(1)} |f(X(s))|^k ds\right) \\ &\leq E\left(\int_{T(0)}^{T(1)} (1 + f(X(s))^{2m}) ds\right) \\ &= E\tau_1 + EY_1(f^{2m}) < \infty \end{aligned}$$

provides the finiteness of the other moment.

By Proposition 2.4, $u(n,k)$ is strongly consistent for $\mu_k(f)$. To state our next result, we shall use the notation $o_p(n^{-1/2})$ to denote the sequence of random variables (r.v.'s) δ_n such that $n^{1/2} \delta_n \Rightarrow 0$ as $n \rightarrow \infty$, where \Rightarrow means weak convergence. The following properties follow easily from our definition and standard results about weak convergence (see p. 92 of CHUNG (1974)):

$$\begin{aligned}
 (2.5) \quad & 1) \text{ if } X_n = o_p(n^{-1/2}) \text{ and } Y_n = o_p(n^{-1/2}), \\
 & \text{then } Z_n = X_n + Y_n = o_p(n^{-1/2}) \\
 & ii) \text{ if } X_n \Rightarrow X \text{ and } Y_n = o_p(n^{-1/2}), \\
 & \text{then } Z_n = X_n \cdot Y_n = o_p(n^{-1/2}).
 \end{aligned}$$

For our next limit result, we will need the following central limit theorem (CLT).

(2.6) PROPOSITION. For $1 \leq k \leq m$, $n^{1/2}(r(n, f^k) - r(f^k)) \Rightarrow \sigma(f^k) N(0,1)$ as $n \rightarrow \infty$, where $\sigma^2(g) = EZ_1^2(g)/(E\tau_1)^2$, $Z_n(g) \equiv Y_n(g) - r(g)\tau_n$, and $N(0,1)$ is a mean zero normal r.v. with unit variance.

Proposition 2.6 is well known in the regenerative simulation literature (see, for example, Iglehart (1978), it is a CLT for estimators of the uncentered steady-state moments of $f(X(\cdot))$.

(2.7) PROPOSITION. $u(n,k) = r(n, f_k) + o_p(n^{-1/2})$ for $1 \leq k \leq m$, where $f_k(\cdot) = f_c^k(\cdot) - k\mu_{k-1}(f) f_c(\cdot)$.

PROOF. Observe that

$$(2.8) \quad u(n, k) = \sum_{j=0}^k \binom{k}{j} r(n, f^j) (-1)^{k-j} r(f)^{k-j} \\ + \sum_{j=0}^{k-1} \binom{k}{j} r(n, f^j) (-1)^{k-j} (r(n, f)^{k-j} - r(f)^{k-j}) .$$

Evidently, for $0 \leq p \leq k$,

$$(2.9) \quad n^{1/2} (r(n, f)^k - r(f)^k) r(n, f^p) \\ = n^{1/2} (r(n, f) - r(f)) \cdot \left(\sum_{j=0}^{k-1} r(n, f)^j r(f)^{k-1-j} \right) r(n, f^p) .$$

From the proof of Proposition 2.4,

$$(2.10) \quad \left(\sum_{j=0}^{k-1} r(n, f)^j r(f)^{k-1-j} \right) \cdot r(n, f^p) \rightarrow k r(f)^{k-1} \cdot r(f^p) \quad \text{a.s.}$$

as $n \rightarrow \infty$, so that (2.9) and (2.10) together imply that

$$(2.11) \quad (r(n, f)^k - r(f)^k) \cdot r(n, f^p) \\ = (r(n, f) - r(f)) \cdot k r(f)^{k-1} \cdot r(f^p) + o_p(n^{-1/2}) .$$

By noting that the first sum on the right-hand side of (2.8) is $r(n, f_c^k)$, we can combine (2.8) and (2.11) to obtain

$$u(n,k) = r(n, f_c^k) + (r(n,f) - r(f))$$

$$\cdot \sum_{j=0}^{k-1} \binom{k}{j} r(f^j) (-1)^{k-j} (k-j) r(f)^{k-j-1} + o_p(n^{-1/2})$$

$$= r(n, f_c^k) - r(n, f_c) \cdot k$$

$$\cdot \sum_{j=0}^{k-1} \binom{k-1}{j} r(f^j) (-1)^{k-1-j} r(f)^{k-j-1} + o_p(n^{-1/2})$$

$$= r(n, f_k) + o_p(n^{-1/2}) .$$

Let $U(n) = (u(n,1), \dots, u(n,m))^T$ and $\mu = (\mu_1(f), \dots, \mu_m(f))^T$.

We take all vectors as column vectors and T denotes transpose.

(2.12) THEOREM. $n^{1/2}(U(n) - \mu) \Rightarrow N(0, C(f))$ where $N(0, C(f))$

is a multivariate normal r.v. with mean vector 0 , and covariance matrix $C(f)$ with elements given by

$$C_{ij}(f) = EZ_1(f_i) Z_1(f_j) / (E\tau_1)^2 .$$

PROOF. Let a be a arbitrary column vector in R^m , and note that Proposition 2.7 implies that

$$\begin{aligned}
& \mathbf{a}^T (U(n) - \mu) \\
&= \sum_{i=1}^m a_i (r(n, f_i) - \mu_i(f)) + o_p(n^{-1/2}) \\
&= \sum_{i=1}^m a_i \bar{Z}_n(f_i) / \bar{\tau}_n + o_p(n^{-1/2}),
\end{aligned}$$

where

$$\bar{Z}_n(g) \equiv n^{-1} \sum_{k=1}^n Z_k(g), \quad \text{for } g : S \rightarrow \mathbb{R}.$$

In the second equality we have used the fact that $\mu_i(f) = r(f_i)$. Standard arguments then show that

$$n^{1/2}(\mathbf{a}^T (U(n) - \mu)) \rightarrow (\mathbf{a}^T \mathbf{C}(f) \mathbf{a})^{1/2} N(0,1)$$

as $n \rightarrow \infty$; the Cramér-Wold device (BILLINGSLEY (1968), p. 48) then yields the desired theorem.

Theorem 2.12 shows that our estimators have an asymptotically normal distribution; it is, of course, a limit theorem expressed in terms of an index n corresponding to the number of regenerative cycles simulated. However, in certain settings, it is more natural to express limit theorems in terms of t , the amount of time that the process \mathbf{X} has been simulated. Then, $N(t) = \max\{n : T(n) \leq t\}$ is the number of regenerative cycles completed by time t . Set

$$u_t(k) = \begin{cases} u(N(t), k); & N(t) \geq 1 \\ 0; & N(t) < 1, \end{cases}$$

and

$$U_t = (u_t(1), \dots, u_t(m)).$$

(2.13) PROPOSITION. $u_t(k) \rightarrow \mu_k(f)$ a.s. as $t \rightarrow \infty$, for $1 \leq k \leq 2m$.

This follows immediately from Proposition 2.4 and the fact that $N(t) \rightarrow \infty$ a.s. as $t \rightarrow \infty$.

(2.14) THEOREM. $t^{1/2}(U_t - \mu) \Rightarrow N(0, C^*(f))$ as $t \rightarrow \infty$, where $C^*(f)$ is given by

$$C_{ij}^*(f) = EZ_1(f_i) Z_1(f_j) / E\tau_1.$$

The proof of Theorem 2.14 is based on making a "random time change" by substituting the process $N(t)$ for the parameter n in Theorem 2.12, giving

$$N(t)^{1/2}(U(N(t)) - \mu) \approx N(0, C(f));$$

since $N(t) \approx t/E\tau_1$, we obtain the result. For a rigorous proof of a more general result, see Theorem 3.10 of GLYNN and IGLEHART (1985).

Specializing our results to the steady-state variance, we observe that the estimator $u(n,2)$ ($u_t(2)$) is asymptotically normal with limiting variance given by $EZ_1(f_2)^2 / (E\tau_1)^2 (EZ_1(f_2)^2 / E\tau_1)$. Since $EZ_1(f_2)^2 = EZ_1(f_2^2)^2$, it follows that the asymptotic variability of our steady-state

variance estimator is unaffected by having to estimate $r(f)$. (Note that $EZ_1(f_c^2)^2 / (E\tau_1)^2$ is the variance of the limiting normal r.v. which approximates $r(n, f_c^2)$.) For higher-order central moments, however, the variances will generally differ.

3. CONFIDENCE INTERVAL GENERATION

In this section, we use our limit theorems of Section 2 to construct confidence intervals for the k 'th central moment. To accomplish this, we need consistent estimators for the covariance matrices $C(f)$ and $C^*(f)$. Set

$$A_k(n, i, j) = \int_{T(k-1)}^{T(k)} (f(X(s)) - r(n, f))^i ds \\ \cdot \int_{T(k-1)}^{T(k)} (f(X(u)) - r(n, f))^j du ,$$

and

$$\bar{A}_n(i, j) = \frac{1}{n} \sum_{k=1}^n A_k(n, i, j) .$$

(3.1) PROPOSITION. For $1 \leq i, j \leq m$,

$$\bar{A}_n(i, j) \rightarrow EY_1(f_c^i) Y_1(f_c^j) \quad \text{a.s. as } n \rightarrow \infty .$$

PROOF. Note that

$$\begin{aligned}
\bar{A}_n(i,j) &= \sum_{k=0}^i \sum_{l=0}^j \binom{i}{k} \binom{j}{l} (-1)^{k+l} r(n,f)^{k+l} \\
&\quad \cdot \frac{1}{n} \sum_{p=1}^n Y_p(f^{i-k}) Y_p(f^{j-l}) \tau_p^{k+l} \\
&\rightarrow \sum_{k=0}^i \sum_{l=0}^j \binom{i}{k} \binom{j}{l} (-1)^{k+l} r(f)^{k+l} \\
&\quad \cdot EY_1(f^{i-k}) Y_1(f^{j-l}) \tau_1^{k+l} \\
&= EY_1(f_c^i) Y_1(f_c^j)
\end{aligned}$$

a.s. as $n \rightarrow \infty$, by the strong law of large numbers.

Let

$$\begin{aligned}
C(n,i,j) &= \frac{1}{-2\tau_n} \{ \bar{A}_n(i,j) - ju(n, j-1) \bar{A}_n(i,1) - u(n,j) \bar{A}_n(i,0) \\
&\quad - iu(n, i-1) \bar{A}_n(1,j) \\
&\quad + iju(n, i-1) u(n, j-1) \bar{A}_n(1,1) \\
&\quad + iu(n, i-1) u(n,j) \bar{A}_n(1,0) \\
&\quad - u(n,i) \bar{A}_n(0,j) \\
&\quad + ju(n,i) u(n,j-1) \bar{A}_n(0,1) \\
&\quad + u(n,i) u(n,j) \bar{A}_n(0,0) \} .
\end{aligned}$$

Propositions 2.4 and 3.1 together yield the following result.

(3.2) PROPOSITION. For $1 \leq i, j \leq m$, $C(n, i, j) \rightarrow C_{ij}(f)$ a.s. as $n \rightarrow \infty$.

Application of the converging-together lemma (BILLINGSLEY (1968), p. 5) to Theorem 2.12 and Proposition 3.2 shows that if $C_{kk}(f) > 0$ ($1 \leq k \leq m$), then

$$\left[u(n, k) - z(\delta) \frac{C_{kk}(f)^{1/2}}{n^{1/2}}, u(n, k) + z(\delta) \frac{C_{kk}(f)^{1/2}}{n^{1/2}} \right]$$

is an asymptotic $100(1-\delta)\%$ confidence interval for $\mu_k(f)$, where $z(\delta)$ solves $P\{N(0,1) \leq z(\delta)\} = 1 - \delta/2$.

A similar confidence interval can be based on simulation of X to time t . Set

$$C_t(i, j) = \begin{cases} C(N(t), i, j) \bar{\tau}_{N(t)} & ; \quad N(t) \leq 1 \\ 0 & ; \quad N(t) < 1 \end{cases}$$

(3.4) PROPOSITION. For $1 \leq i, j \leq m$, $C_t(i, j) \rightarrow C_{ij}^*(f)$ a.s. as $t \rightarrow \infty$.

This result is an immediate consequence of Proposition 3.2, and leads to the following asymptotic $100(1-\delta)\%$ confidence interval for $\mu_k(f)$ (assuming $C_{kk}(f) > 0$, $1 \leq k \leq m$):

$$(3.5) \quad \left[u_t(k) - z(\delta) \frac{C_{kk}^*(f)^{1/2}}{t^{1/2}}, u_t(k) + z(\delta) \frac{C_{kk}^*(f)^{1/2}}{t^{1/2}} \right]$$

For the steady-state variance $v(f) = \mu_2(f)$, we can use our knowledge that $\mu_1(f) = 0$ to obtain a simpler family of estimates for $C_{22}(f)$ and $C_{22}^*(f)$. Note that

$$\begin{aligned} (3.6) \quad EZ_1(f_2)^2 &= EZ_1(f_c^2)^2 \\ &= EY_1(f_c^2)^2 - 2v(f) EY_1(f_c)\tau_1 + v(f)^2 E\tau_1^2. \end{aligned}$$

Set

$$C^0(n) = \frac{1}{\tau_n^2} \{ \bar{A}_n(2,2) - 2u(n,2) \bar{A}_n(2,0) + u(n,2)^2 \bar{A}_n(0,0) \}$$

$$C_t^0 = \begin{cases} C^0(N(t)) \bar{\tau}_{N(t)} & ; \quad N(t) \geq 1 \\ 0 & ; \quad N(t) < 1 \end{cases}$$

Propositions 2.4 and 3.1 show that $C^0(n) \rightarrow C_{1j}(f)$ a.s. as $n \rightarrow \infty$ and $C_t^0 \rightarrow C_{1j}(f)$ a.s. as $t \rightarrow \infty$, provided $n \geq 2$. Thus, if $C_{22}(f) > 0$, the following intervals are $100(1-\delta)\%$ asymptotic confidence intervals for $v(f)$:

$$(3.7) \quad \left[u(2,n) - z(\delta) \frac{C^0(n)^{1/2}}{n^{1/2}}, u(2,n) + z(\delta) \frac{C^0(n)^{1/2}}{n^{1/2}} \right]$$

$$(3.8) \quad \left[u_t(2) - z(\delta) \frac{(C_t^0)^{1/2}}{t^{1/2}}, u_t(2) + z(\delta) \frac{(C_t^0)^{1/2}}{t^{1/2}} \right].$$

4. NUMERICAL RESULTS

To illustrate the results obtained in the previous section we have simulated three models: the waiting time process in the M/M/1 queue ($\rho = 0.5$), an (s,S) inventory model, and the classical repairman model. For all three models we selected f to be the identity function ($f(x) = x$) and $k = 2$, so that our goal was to estimate the variance of the steady-state distribution.

(4.1) **EXAMPLE. M/M/1 Queue.** This model is the single server queue with Poisson arrivals and exponential service times. We simulate the waiting time process $W = \{W_n : n \geq 0\}$, where W_n is the waiting time (exclusive of service time) of the n th customer. Our simulations were carried out for arrival rate $\lambda = 5$ and service rate $\mu = 10$, and hence the traffic intensity $\rho = 0.5$. This guarantees that $W_n \Rightarrow W$ as $n \rightarrow \infty$.

Regenerative cycles begin at those values of n for which $W_n = 0$. The quantity being estimated here is $\sigma^2\{W\} = 3.0$. We did 50 replications of 5000 cycles each. The sample mean of the 50 point estimates, $u(2,5000)$, was 3.0417, and the sample mean of the 50 point estimates of $C_{22}(f)^{1/2}$, $C^0(5000)^{1/2}$, was 13.2782. As a result, the sample mean of the 90% confidence intervals was [2.7328, 3.3506]. The coverage probability was 58%.

(4.2) **EXAMPLE. (s,S) Inventory Model.** This model is a periodic review inventory model with a stationary (s,S) ordering policy. An (s,S) policy is characterized by two positive integers: s and S with $s < S$. If the

amount of inventory on hand plus on order is less than s , order to bring the sum up to S . If the inventory is greater than or equal to s , do not order. Let X_n denote the level of inventory on hand plus on order in period n immediately after the ordering decision. If d_n denotes the demand in period n , then

$$X_{n+1} = \begin{cases} X_n - d_n, & d_n \leq X_n - s \\ S, & \text{otherwise.} \end{cases}$$

We assume that $s \leq X_0 \leq S$. The state space of $\{X_n : n \geq 0\}$ is $\{s, s+1, \dots, S\}$. For this example we have selected $s = 6$, $S = 10$, and

$$P\{d_n = j\} = \begin{cases} 3/8, & j = 0 \\ 1/4, & j = 1 \\ 3/16, & j = 2 \\ 1/8, & j = 3 \\ 1/16, & j = 4 \end{cases}$$

Again we simulate to estimate $\sigma^2\{X\} = 2.3333$. Using $i = 10$ as the regenerative state we run 50 replications of 1000 cycles each. The sample mean of the 50 point estimates, $u(2,1000)$, was 2.3352 and the sample mean of the 50 point estimates for $C_{22}(f)^{1/2}$, $C^0(1000)^{1/2}$, was 1.2396. The sample mean of the resulting 50 90% confidence intervals was [2.2708, 2.3997]. The coverage probability was 94%.

(4.3) **EXAMPLE. Classical Repairman Model.** This model is a continuous time Markov chain with $X(t)$ denoting the number of failed units undergoing or waiting for service at the repair facility at time t . We have

$m + n$ identical machines each with an exponential failure time with failure rate λ . At most n of the units operate at one time, the other m being thought of as spares. When a unit fails, it is sent to a repair facility consisting of s repairmen (servers) having exponential repair (service) times with repair rate μ . With these assumptions $\{X(t) : t \geq 0\}$ is a birth-death process with state space $\{0, 1, \dots, m + n\}$, birth parameters $\lambda_i = (n - [i - m]^+) \lambda$, and death parameters $\mu_i = \mu \cdot \min(i, s)$. For this example we have used $n = 10$, $m = 4$, $\lambda = 1$, $\mu = 4$, and $s = 3$. Again we are interested in estimating the steady state variance, $\sigma^2\{X\} = 5.231$. We ran 50 replications of 1000 cycles each with the regenerative state taken to be $i = 2$. The sample mean of our 50 point estimates, $u(2, 1000)$, was 5.1916, and the sample mean of the 50 point estimates of $C_{22}(f)^{1/2}$, $C^0(5000)^{1/2}$, was 11.5562. The sample mean of the 50 90% confidence intervals was [4.5905, 5.7927]. The coverage probability was 74%.

REFERENCES

- BILLINGSLEY, P. (1968). Convergence of Probability Measures. John Wiley, New York.
- CHUNG, K.L. (1974). A Course in Probability Theory. Academic Press, New York.
- GLYNN, P.W. and D.L. IGLEHART (1985). The joint limit distribution of the sample mean and regenerative variance estimator. Forthcoming technical report, Department of Operations Research, Stanford University, Stanford, CA.
- HEYMAN, D.P. and M.J. SOBEL (1982). Stochastic Models in Operations Research, Volume 1. McGraw-Hill, New York.
- IGLEHART, D.L. (1978). The regenerative method for simulation analysis. In Current Trends in Programming Methodology -- Software Modeling (K.M. Chandy and R.T. Yeh, editors). Prentice-Hall, Englewood Cliffs, N.J.

END

FILMED

1-86

DTIC